

RECLAMATION OF SODIC SOILS BY DIFFERENT DOSES OF GYPSUM TO ENHANCE GRAIN YIELD AND QUALITY OF RICE (*ORYZA SATIVA* L.) IN SULTANPUR DISTRICT, U.P.

Trilok Nath Rai², Kedar Nath Rai¹, Sanjeev Kumar Rai³, Pankaj Nautiyal⁴ & Traloki Singh⁵
^{1&3}Department of Soil Science & Agricultural Chemistry, Kamla Nehru Institute of Physical and Social Sciences, Sultanpur, Uttar Pradesh, India
^{2,4&5}Department of Soil Science & Agricultural Chemistry, Krishi Vigyan Kendra (ICAR-CSSRI), Dhikunni, Sandila, Hardoi, Uttar Pradesh, India

Salinity is a global problem reducing plant growth and productivity worldwide, and affects about 7% of the world's total land area (Flower *et al.* 1997). In the world about 1200 million hectare (Mha) of land is affected with salinity and, therefore, poses a challenging task of taking up agriculture and enhancing productivity in these areas (Poonamperuma 1984; Tanji 1990). Some of the most severe problem in soil salinity are reported in arid and semiarid regions of the world (Pesarrakli 1999), where limited rainfall, high evapo-transpiration and high temperature play an important role in increasing the salt concentration in the root zone. There could be two possible ways to overcome salt stress (i) either improve the soil environment for the normal plant growth through leaching of salts from the profile through chemical amendment like gypsum ($\text{CaSO}_4 \cdot 2\text{H}_2\text{O}$) (Meri 1984; Al-Nabulsi 2001) or (ii) improve the plant itself which can be grown in that environment (Jardat *et al.* 2004; El-Hendawy *et al.* 2005). The first approach is resource costly and the second is based on the development of salt tolerant varieties. There could be a third approach, *i.e.* hybrid approach, based on exploitation of synergies between gypsum and salt tolerant varieties (Singh *et al.* 2009). Although million of hectares of salt-affected soils are potentially suitable for crop production with appropriate improvement measures, they are left uncultivated or are grown with crops with very low yields because of salinity problem.

Rice (*Oryza sativa* L.) is one of the most important cereal crops grown in wide range of climatic zones to nourish the mankind. Rice varieties suited to normal conditions may rarely or mostly not adapt under salt stress conditions. Few screening studies have been reported based on stability of rice genotypes across sodicity stress and non-stress environments (Zapata *et al.* 1991; Shylaraj *et al.* 1994) while no such study has been reported for sodicity stress. Sodicity being a specific agro-edaphic environment spread over about 3.8 Mha area of the country (NRSA and Associates 1996) for which gypsum ($\text{CaSO}_4 \cdot 2\text{H}_2\text{O}$) is generally recommended for the reclamation. Cultivation of rice is generally recommended as first crop after application of gypsum because continuous submergence improves soil properties. Gypsum application markedly decreased the soil pH, exchangeable sodium percentage (ESP) and significantly increased the yield and nutrient uptake by rice (Singh *et al.* 2008).

More than 70% farmers in the region of sodic soils belong to small and marginal categories and the initial cost of reclamation is beyond the reach of this category of farmers because of heavy

investment on account of gypsum (Dutta *et al.* 1996). Earlier studies revealed that judicious and proper use of gypsum can markedly increase the yield and nutrient uptake of rice. However, little information is available on the combined effect of gypsum and varieties on soil amelioration, grain yield and quality of rice. Therefore, given the importance of gypsum on sodic soil amelioration and grain yield of rice, it is necessary to know the extent of benefit a rice variety, which can confer to increase yield and associated components. Therefore, a three years study was undertaken to have a detailed account of the response of rice varieties and soil reclamation to gypsum.

MATERIALS AND METHODS

Site Characterization

A pot experiment to find out the response of gypsum on growth, yield and quality of rice and soil health was conducted in barren sodic soil at Motigarpur Sultanpur, Uttar Pradesh in controlled conditions for 3 years from 2017 to 2019. The mean annual rainfall is 855 mm and more than 85% generally occurs during monsoon season (July to September). Mean annual soil temperature varies from 18.25°C during winter to 32.5°C during summer. The soil of the experimental field was highly sodic with pH (1: 2, soil:Water) 10.5, electrical conductivity (EC) 2.42 dS m⁻¹ and ESP 89, having low organic carbon (0.80 g kg⁻¹) and available N (94kg ha⁻¹), medium available P (25kg ha⁻¹) and rich in available (388.8kg ha⁻¹) at 0-15 cm soil depth. The GR value of the experimental soil determined by Schoonover (1952) method at the time of initiating the trials (2017) was 30.8 t ha⁻¹.

Experimental Details

A four-times replicated experiment was laid out in a split plot design with four gypsum levels (control, 13% GR, 23% GR and 52% GR) as main plot treatment and two varieties 'UD 3' and 'Pant 10' as sub-plot treatment in an elementary pot of 50m² size. Both the varieties were planted randomly in each gypsum pot covering 25 m² area. As per treatments, gypsum was incorporated once in surface soil upto 10 cm depth in the month of June and about 10 cm water was ponded in the pots for 10 days to displace the reaction products of Ca-Na exchange down the root zone. 45 days-old seedlings of two rice varieties 'UD 3' and 'Pant10' were transplanted at 20 cm row to row and 15 cm plant to plant spacing during second week of July every year. The recommended doses for sodic soils of N (165 kg ha⁻¹) and zinc sulphate (ZnSO₄) (50 kg ha⁻¹) were applied uniformly in all the treatments. Basal fertilizer schedule consisted of half dose of N and full dose of zinc sulphate (50 kg ha⁻¹) were applied uniformly in all the treatments. The remaining half the dose of was applied in two equal splits at 35 and 65 days after transplanting. Pots received identical cultural treatments in terms of ploughing, cultivation, transplanting method and disease control, *etc.*

Plant Growth and Yield Parameters

Five hills in each pot were randomly selected and tagged for recording growth parameters like plant height, numbers of effective tillers/hill, number of leaves/hill, leaf weight and leaf area index. Dry matter accumulation was recorded at 35 days interval from 3 hills/pot harvested from outside the net pot area. Days to 50% flowering was recorded from the number of panicles emerged in a unit area. The net plots area (5 m² for each variety) was harvested and the biological yield was recorded. Yield attributes *viz.*, length of panicle, grains/panicle and 1000grain weight were recorded from tagged plants. The grain and straw yields were recorded after threshing, cleaning and drying of produce and straw yield was obtained by subtracting grain yield from total biomass yield. The benefit:cost (B:C) ratio was computed on the basis of prevailing market price of produce and local cost of inputs.

Grain Quality Analysis

Nitrogen content in grain was analyzed following Kjeldahl method using block digestion and steam distillation method (Foss Analytical 2003). Length: width ratio of grain with and without husk, length:width ratio of cooked rice, brown rice recovery, milled rice recovery, head rice recovery and gel consistency at 35 and 65 min after cooking were analyzed as per standard methods.

Soil Analysis

Soil samples (0-15 cm) were taken every year after harvesting of rice crop and analyzed to monitor the changes in soil properties due to gypsum doses and rice varieties. Air-dried soil samples were ground to pass through 2-mm sieve and analyzed for pH and EC using a glass electrode. The OC content was determined by Walkley and Black method (Jackson 1973). Available N was determined by distillation of soil with KMnO_4 and NaOH (Subbiah and Asija 1956). Exchangeable Na^+ percentage (ESP) was calculated by the formula $\text{ESP} = [\text{exchangeable } \text{Na}^+ (\text{cmol}(\text{p}^+) \text{ kg}^{-1}) \times 100/\text{CEC} (\text{cmol}(\text{p}^+) \text{ kg}^{-1})]$ (Richards, 1954).

Statistical Analysis

A two-way analysis of variance (ANOVA) was used to test the effect of gypsum doses and varieties as well as their interaction on growth, yield and quality of rice. The data were analyzed using the statistical package MSTATC. The differences between the gypsum mean effects were compared using the least significant differences (*LSD*) at the probability level $P= 0.05$.

RESULTS AND DISCUSSION

Effect of Gypsum on Soil Reclamation

After three years of rice, the pH of surface soil (0-15 cm) treated with gypsum at 52% GR declined from 10.5 to 9.10 whereas, it decreased to 9.34 in the treatment receiving gypsum at 23% GR for the 'Pant10' variety (Table1). In control plot, it declined to 10.01. Growing the 'UD3' variety and applying 23 and 52%GR doses of gypsum reduced the soil Ph to 9.21 and 9.00 whereas, growing the 'Pant 10' variety declined it to 9.34 and 9.10, respectively. There was no significant difference in soil pH due to varieties. It may be because sodic soils contain measurable amount of NaHCO_3 and Na_2CO_3 which under normal conditions react with added gypsum ($\text{CaSO}_4 \cdot 2\text{H}_2\text{O}$) to neutralize and precipitate soluble Ca before it can be used to replace exchangeable Na.

The ESP of the surface soil declined from 89 to 45.4 and 45.7 at 23% GR for UD 3 and Pant 10, respectively; however, it reduced to 32.0 and 34.4 at 52% GR after three years. There was no significant difference in ESP due to varieties but combined effect of gypsum and varieties plays a significant role in ESP levels (Singh *et al.* 2008). One possible reason may be because of cultivation of rice crop which has capacity to solubilize soil native CaCO_3 to further reduce soil exchangeable sodium. Chhabra and Abrol (1977) have also reported the changes in ESP and improvement in soil properties with gypsum doses and cultivation of rice. Combined effect of gypsum and rice cultivars has generated more organic acids, which mobilize the soil calcium to leach down the salts from the root zone. Addition of gypsum increased the OC content of the soil. After three years of rice with 'UD 3' and 'Pant 10' at 23 and 52% GR levels, the OC content of the surface soil increased significantly to 1.18, 1.04, 1.00 and 1.20 g kg^{-1} , respectively over the initial value of 0.80 g kg^{-1} . After one year of study *i.e.* after first year, available N content with 13, 23 and 52% GR increased by 8.4, 33.4 and 42 percent over the control for the 'UD 3' variety. Available N content after three years increased significantly to

190 kg ha⁻¹ with 52% GR which is about 72.6 per cent over the control. However, difference between the varieties in this character was not significant. There may be one possible reason that enhanced organic matter in soil and total biomass generated with respect to gypsum levels and growing of rice which indirectly facilitates the removal of exchangeable Na⁺ by increasing the cross-sectional area of conducting pores, resulting in increased permeability and generation of more organic acids that mobilize the soil calcium. Rice roots provide channels for the movement of water, which increased permeability (Chhabra and Abrol 1977).

Effect of Gypsum on Plant Growth

Plant height was significantly ($P \leq 0.05$) increased by increasing doses of gypsum. The increment in plant height was 50.3, 58.1 and 67.8 per cent at 13, 23 and 52% GR doses compared with the control treatment, respectively. A non-significant interaction between gypsum and varieties on plant height was observed (Table 2). The increase in plant height in response to gypsum levels is due to reduction in soil pH and exchangeable sodium and increase in OC and available N (Table 1) which might have enhanced more leaf area resulting into higher photo assimilates and thereby resulted in more dry matter accumulation. The growth reduction in sodic soils could also be the result of toxic effects related to the accumulation of Na⁺ ions (Ehert *et al.* 1990; Brugnoli and Lauter 1991; Saneoka *et al.* 1999; Akhtar *et al.* 2001).

Number of effective tillers/hill at harvests (105- 125 days after transplanting) increased significantly with increase in doses of gypsum (Table2). Gypsum at 50% GR produced the maximum number of effective tillers/hill (9.86) but the difference between 23 and 52% GR levels was not statistically significant. However, there was no significant difference between varieties in this character. Numbers of green leaves/hill were significantly higher with 52% GR but statistically at par with that of 23% GR. A similar pattern was also registered with respect to leaf weight/ hill. The increase in leaf count as well as leaf weight/ hill with increasing levels of gypsum may be because of reduction in soil pH and exchangeable sodium and increasing OC content and available N in soil system (Table 1) which, in turn, increased photosynthesis and assimilates the photosynthates. Singh *et al.* (1983) have reported that addition of gypsum increased leaf count, leaf weight as well as leaf area which increased nutrient availability in the plants as a result of better root development and increasing N use efficiency. Application of gypsum at 52% GR recorded maximum leaf area index (2.21) while the minimum (1.22) was observed in the control plot. It might be due to improved soil health with increasing levels of gypsum, higher nutrient availability and enhanced plant growth.

The data presented in table1 revealed a statistically significant increase in dry matter due to increasing levels of gypsum. Significantly higher dry matter accumulation (61.8 g/hill) was obtained from 52% GR at harvest of crop. These results were statistically at par with that of 23% GR because of almost similar vegetative growth due to non-significant difference in soil pH, OC and available N status of the soil (Table1). The higher dry matter yield from higher levels of gypsum could be due to more availability of N to the plants (Obrejanu and Sandhu 1971). These differences were statistically significant over the control for which the lowest dry matter accumulation (19.7 g/hill) was obtained.

Days to 50% flowering and days to maturity were also affected with increasing levels of gypsum. 50% flowering in control plot was recorded about 12- 20 days earlier than the gypsum treated plots. It might be due to physiological stress in the plant. Similar trend was also observed in days to

maturity. This result is consistent with the findings of Kings bury and Epstein (1986) and Houshmand *et al.* (2005). All the growth parameters of 'UD 3' were significantly superior over the 'Pant 10'.

Effect of Gypsum on Yield Attributes

Panicle length in rice increased with increase in gypsum rates (Table 3) up to 52% GR level but significant difference in this character was recorded only up to 23% GR level. However, a non significant effect of variety×gypsum interaction was observed. Number of grains/panicle differed significantly with the levels of gypsum. A significant interaction was also exhibited between varieties × gypsum levels (Table 3). Again, in terms of test weight, 23 and 52% GR value were at par between themselves but significantly superior over control and 13% GR levels with a non significant variety × gypsum interaction. Variety 'UD 3' gave significantly higher panicle length over the 'Pant 10'. This might be due to more root proliferation and increasing N use efficiency of UD 3 in the gypsum treated plots (Kumar *et al.* 1994). Similarly, the test weight of 'UD 3' was significantly higher than the traditional variety 'Pant 10' because of bolder grain size (Table 3). It appears that the application of gypsum, which contains sulphur, increased the protein percentage, which in turn increased the grain weight (Kadamdhad *et al.* 1996).

Effect of Gypsum on Grain, Straw and Biological Yields

The grain yield data (Tables 3 and 4) indicate a positive response to gypsum application. The pooled data revealed that application of gypsum at 52% GR gave the maximum grain yield of rice (4.75 t ha⁻¹), which was statistically at par with that of 23% GR (4.35 t ha⁻¹) understandably because of the similar trend in yield attributing characters like length of panicle, number of grains/panicle and test weight. Chhabra *et al.* (1989) have also reported non- significant difference in rice grain yield between 23 and 52% GR levels. Similarly, Arora *et al.* (2015) recorded highest grain and straw yield of CSR36 paddy with gypsum @50%GR compared to 25%GR in sodic soils. The decrease in yield under control, might have occurred due to retarded growth of the plants as a result of the low uptake of water and nutrients as well as the ion-toxic effects of Na⁺ (Yeo and Flower 1986; Flower *et al.* 1990; Akhtar *et al.* 2001). The variety 'UD 3' gave significantly higher grain yield than 'Pant 10' and a significant interaction between varieties × gypsum on grain yield was observed. It indicates that varieties responded to sodicity levels differently. The mean increase in grain yield of 'UD3' was 21 per cent over traditional variety 'Pant 10'. Sridhar *et al.* (1985) reported higher response to gypsum in respect of grain yield which might be due to readily available S & Ca in gypsum. Straw yield with 52% GR was significantly higher over control and 13% GR levels because of higher plant height (Table1) and dry matter accumulation (Table 2) but it was at par with that of 23% GR. It was observed that 52% GR gave the maximum straw yield (10.2 t ha⁻¹) while the lowest straw yield (2.70 t ha⁻¹) was obtained from the control treatment. Cultivar 'UD3' gave significantly higher straw yield over 'Pant10' but the interaction between varieties× gypsum was non-significant. Similar trend was also observed in biological yield. The mean increase in straw and biological yields of 'UD 3' was 20.1 and 20.2 per cent, respectively, over the traditional variety 'Pant 10'. The interaction between varieties × gypsum for straw and biological yields was non-significant.

Effect of Gypsum on Harvest Index and B:C Ratio

The harvest index was significantly increased by increasing levels of gypsum. Application of gypsum at 52% GR recorded the maximal harvest index which was statistically at par with that of

23%GR because of similar trend in biological and grain yields (Table3). Variety 'UD3' was significantly superior over pant 10 in this character a non-significant interaction between gypsum and varieties on harvest index was observed. Numerically highest benefit: cost (B:C) ratio (1.78 based on pooled data) was recorded with 52% GR; however, it was statistically at par with the application of gypsum at 23% GR (1.76) (Table 3). The benefit: cost ratio observed from variety 'UD 3' was significantly higher over traditional variety 'Pant10'. The interaction between gypsum and varieties on B:C ratio was non-significant.

Effect of Gypsum on Grain Quality

Nitrogen content in rice grain is markedly influenced by the sodicity levels (Table 5). The significantly lower N content for control could be due to high volatilization losses of N at high pH resulting in slow transformation of N from amide to ammonia and nitrate and its less availability to the plants, resulting in less N uptake. Nitant and Bhumbra (1974) reported that complete hydrolysis of urea, the most commonly used nitrogenous fertilizer, is delayed due to higher soil Ph than that in soil of low pH. The reduced N content in grain in control plots could be due to the impaired availability of N at higher pH, which reduced nitrogen uptake in the plants. High pH and high amount of CaCO₃ also favor volatilization losses of applied N (Rao and Batra 1983).

Application of gypsum improved the quality of rice in terms of rice grain length and width, grain length: breadth ratio with and without husk, length: breadth ratio of cooked rice, brown rice recovery, milled rice recovery, head rice recovery and gel consistency (Table5). Data revealed that application of gypsum at 52% GR, recorded maximum rice grain length and breadth of 9.30 and 2.77 mm, respectively. These results were statistically at par with that of 23% GR giving corresponding figures of 9.20 & 2.77 mm. The significantly lowest length (8.98 mm) and breadth (2.69 mm) of grain were recorded with the control. The rice recoveries in terms of brown rice, milled rice and head rice were also increased with increasing levels of gypsum. Head rice recovery was higher for 52% GR over the control, 13 %GR and 23% GR. However, milled rice recovery and brown rice recovery were statistically at par with that of 23% GR. This might be due to the increase in availability of S and Ca from applied gypsum to the plant and its subsequent utilization for grain development. Variety 'UD 3' was significantly superior over 'Pant 10' in terms of grain N, grain length, grain breadth with husk, length: width ratio with husk, length: breadth ratio of cooked rice, milled rice recovery, head rice recovery and gel consistency after 62 min. The interaction effect of gypsum × varieties on grain quality was non-significant in most of the rice recovery quality parameters (Table 5).

Table1. Improvement in soil properties due to the combined effect of gypsum and rice varieties

| Soil properties | Control | 13%GR | 23%GR | 52%GR | LSD | | | | |
|------------------------|---------|-------|-------|-------|-------|-------|-------|-------|------|
| First year | | | | | | | | | |
| pH(1:2) | 10.18 | 10.24 | 9.95 | 9.98 | 9.50 | 9.58 | 9.22 | 9.40 | 0.48 |
| EC(dSm ⁻¹) | 0.97 | 1.01 | 0.64 | 0.88 | 0.67 | 0.69 | 0.65 | 0.69 | 0.08 |
| ESP | 82.1 | 87.21 | 72.62 | 75.43 | 60.00 | 63.67 | 45.50 | 52.76 | 6.34 |

| | | | | | | | | | |
|--|-------|-------|--------|--------|--------|--------|--------|--------|------|
| OC(gkg ⁻¹) | 0.84 | 0.81 | 1.04 | 0.86 | 1.08 | 0.90 | 1.10 | 1.00 | 0.02 |
| Alkaline-N (kgha ⁻¹) | 96.4 | 97.10 | 104.52 | 100.25 | 128.60 | 124.30 | 146.62 | 140.50 | 8.92 |
| Second year | | | | | | | | | |
| pH(1:2) | 10.01 | 10.04 | 9.61 | 9.75 | 9.30 | 9.40 | 9.12 | 9.22 | 0.33 |
| EC(1:2)(dSm ⁻¹) | 0.82 | 0.82 | 0.63 | 0.74 | 0.59 | 0.63 | 0.53 | 0.67 | 0.04 |
| ESP | 80.0 | 82.6 | 62.5 | 66.6 | 52.6 | 54.8 | 42.6 | 45.4 | 8.12 |
| OC(gkg ⁻¹) | 0.86 | 0.84 | 1.06 | 0.88 | 1.10 | 0.95 | 1.15 | 1.16 | 0.04 |
| AlkalineKMnO ₄ -N(kgha ⁻¹) | 102.3 | 100.2 | 120.5 | 118.4 | 150.2 | 144.6 | 171.4 | 167.5 | 7.68 |
| Third year | | | | | | | | | |
| pH(1:2) | 9.88 | 10.01 | 9.46 | 9.51 | 9.21 | 9.34 | 9.00 | 9.10 | 0.64 |
| EC(1:2)(dSm ⁻¹) | 0.76 | 0.82 | 0.60 | 0.66 | 0.56 | 0.62 | 0.48 | 0.54 | 0.06 |
| ESP | 72.2 | 75.1 | 55.2 | 57.3 | 45.4 | 45.7 | 32.0 | 34.4 | 7.68 |
| OC(gkg ⁻¹) | 0.90 | 0.86 | 1.06 | 0.90 | 1.18 | 1.00 | 1.20 | 1.20 | 0.03 |
| Alkaline KMnO ₄ - N(kgha ⁻¹) | 110.2 | 110.1 | 143.5 | 142.5 | 172.6 | 168.5 | 190.2 | 190.2 | 4.63 |

| | | | | | | |
|------|---------|------|---------|----------|---------|----|
| UD 3 | Pant 10 | UD 3 | Pant 10 | UD 3 | Pant 10 | UD |
| 3 | | | Pant 10 | (P□0.05) | | |

Table 2. Response of growth parameters of rice varieties to gypsum (3 years pooled data)

| Treatments | Plant height (cm) | No. of effective tillers/hill | Dry matter (g/hill) | No. of leaves/ hill | Leaf weight (g/hill) | Leaf area index | Number of panicle/ m ² | Days to 50% flowering | Days to maturity |
|-----------------------------|----------------------|-------------------------------|------------------------|------------------------|-------------------------|--------------------|--------------------------------------|--------------------------|------------------|
| Gypsum levels (% GR) | | | | | | | | | |
| 0 | 62.1 | 3.80 | 19.7 | 33.3 | 22.2 | 1.22 | 95.4 | 90 | 122 |
| 13 | 93.3 | 6.96 | 44.1 | 51.5 | 32.5 | 1.54 | 170.4 | 101 | 132 |
| 23 | 98.2 | 8.16 | 57.6 | 72.2 | 39.6 | 1.88 | 386.5 | 105 | 140 |
| 52 | 104.2 | 9.86 | 61.8 | 82.1 | 43.5 | 2.21 | 435.2 | 108 | 145 |
| <i>SEm</i> ± | 5.24 | 0.34 | 1.67 | 3.12 | 0.02 | 0.021 | 7.10 | 0.87 | 1.46 |

| | | | | | | | | | |
|-----------------------------|------|------|------|------|-------|------|-------|------|------|
| <i>LSD(P</i> □ <i>0.05)</i> | 16.5 | 1.09 | 5.29 | 9.12 | 0.08 | 0.06 | 21.2 | 2.63 | 4.52 |
| Varieties | | | | | | | | | |
| UD 3 | 94.2 | 8.36 | 55.8 | 64.8 | 36.84 | 1.84 | 324.5 | 102 | 135 |
| Pant 10 | 84.6 | 6.02 | 47.9 | 51.4 | 31.43 | 1.62 | 312.2 | 108 | 142 |
| <i>SEm</i> ± | 1.70 | 1.20 | 0.87 | 1.46 | 0.54 | 0.01 | 2.25 | 0.76 | 1.01 |
| <i>LSD(P</i> □ <i>0.05)</i> | 4.70 | 3.60 | 2.69 | 4.52 | 1.64 | 0.04 | 6.74 | 2.12 | 2.63 |
| Interaction | NS | NS | * | * | * | * | * | * | * |

*Significant at the 0.05 probability level, *SEm* ±: Standard error of means , *LSD*: Least significant difference

Table 3. Response of yield attributes, yields, harvest index and benefit:cost ratio of two rice varieties to gypsum (3 years pooled data)

| Treatments | Length of panicle | No.of grains/ panicle | Test weight (g) | Biological yield (tha ⁻¹) | Straw yield (tha ⁻¹) | Grain yield (tha ⁻¹) | Harvest index | B:C ratio |
|-----------------------------|-------------------|-----------------------|-----------------|---------------------------------------|----------------------------------|----------------------------------|---------------|-----------|
| | (cm) | | | | | | | |
| Gypsum levels (% GR) | | | | | | | | |
| 0 | 15.8 | 29.6 | 17.6 | 2.71 | 2.35 | 0.36 | 13.2 | 0.39 |
| 13 | 21.9 | 92.3 | 22.8 | 9.86 | 6.87 | 2.99 | 30.3 | 1.37 |
| 23 | 24.3 | 126.2 | 24.8 | 13.03 | 8.68 | 4.35 | 33.3 | 1.76 |
| 52 | 25.6 | 137.3 | 25.7 | 14.19 | 9.44 | 4.75 | 33.4 | 1.78 |
| <i>SEm</i> ± | 1.06 | 1.76 | 0.67 | 0.42 | 0.32 | 0.16 | 0.007 | 0.02 |
| <i>LSD(P</i> □ <i>0.05)</i> | 3.36 | 5.72 | 1.91 | 1.30 | 1.05 | 0.50 | 0.026 | 0.07 |
| Varieties | | | | | | | | |
| UD 3 | 21.8 | 103.0 | 24.8 | 10.8 | 7.47 | 3.40 | 31.2 | 1.60 |
| Pant 10 | 22.2 | 89.7 | 20.7 | 9.0 | 6.22 | 2.82 | 31.1 | 1.49 |
| <i>SEm</i> ± | 0.53 | 1.54 | 0.60 | 0.42 | 0.16 | 0.07 | 0.004 | 0.02 |
| <i>LSD(P</i> □ <i>0.05)</i> | NS? | 4.71 | 1.65 | 1.23 | 0.51 | 0.23 | 0.017 | 0.06 |
| Interaction | NS | * | NS | NS | NS | * | NS | * |

*Significant at the 0.05 probability level, ns: non-significant at P=0.05.

Table 4. Analysis of variance for the response of plant height, number of effective tillers, dry matter accumulation, days to 50% flowering, days to maturity, length of panicle, test weight, biological, straw and grain yields to gypsum and varieties

| Characters | Replication | Gypsum | Varieties | Gypsum×varieties |
|---|-------------|----------|-----------|------------------|
| (3d.f.) | | (3d.f.) | (1d.f.) | (3d.f.) |
| Plant height | 1.925 | 12.976* | 15.513* | 0.250 |
| No.ofeffective tillers/hill | 1.068 | 55.180* | 55.086* | 1.500 |
| Dry matter accumulation | 0.152 | 128.798* | 37.534* | 4.628* |
| Days to50% flowering | 0.630 | 9.681* | 12.631* | 3.102* |
| Days to maturity | 0.320 | 14.342* | 9.861* | 4.130* |
| Length of panicle | 0.224 | 16.064* | 0.351 | 0.640 |
| Test weight | 1.143 | 22.835* | 22.844* | 0.497 |
| Biological yield | 0.463 | 158.445* | 38.457* | 8.312* |
| Straw yield | 0.625 | 92.250* | 22.957* | 6.851* |
| Grain yield | 0.170 | 471.646* | 101.626* | 5.134* |
| *Significant at the 0.05 probability level. | | | | |

Table 5. Response of grain quality of rice varieties to gypsum

| Treatments | Grain | Grain | Grain | L:B | L:B | L:B | Brown | Milled | Head | Gel | Gel |
|-------------|---------------------------|--------|---------|-------|---------|---------|----------|----------|----------|-----------------|-----------------|
| | N(%) | length | breadth | ratio | ratio | ratioof | rice | rice | rice | consiste ncy | consiste ncy |
| | | with | with | with | without | cooked | recovery | recovery | recovery | after30 | after60 |
| | | husk | husk | husk | husk | rice | (%) | (%) | (%) | minutes | minutes |
| | | (mm) | (mm) | | | | | | | | |
| | | (L) | (B) | | | | | | | | |
| | Gypsum level (%GR) | | | | | | | | | | |
| 0 | 1.14 | 8.98 | 2.69 | 3.33 | 2.81 | 2.45 | 77.95 | 73.16 | 57.80 | 5.50 | 5.26 |
| 13 | 1.25 | 9.09 | 2.73 | 3.32 | 2.82 | 2.47 | 78.02 | 75.10 | 57.80 | 5.82 | 5.36 |
| 23 | 1.32 | 9.20 | 2.77 | 3.32 | 2.85 | 2.51 | 78.26 | 76.00 | 57.92 | 5.83 | 6.40 |
| 52 | 1.35 | 9.30 | 2.77 | 3.35 | 2.85 | 2.62 | 78.87 | 76.60 | 61.67 | 5.85 | 6.65 |
| <i>SEm±</i> | 0.10 | 0.012 | 0.013 | 0.01 | 0.003 | 0.006 | 0.56 | 0.48 | 0.51 | 0.006 | 0.012 |

| | | | | | | | | | | | |
|---------------------------------|------|-------|-------|------|-------|------|-------|-------|-------|-------|-------|
| <i>LSD</i> ($P \leq 0.05$) | 0.03 | 0.04 | 0.04 | 0.03 | 0.01 | 0.02 | 1.75 | 1.44 | 1.56 | 0.02 | 0.04 |
| Varieties | | | | | | | | | | | |
| UD 3 | 1.26 | 9.15 | 2.73 | 3.35 | 2.82 | 2.46 | 78.50 | 75.20 | 57.20 | 5.84 | 5.75 |
| Pant 10 | 1.28 | 9.16 | 2.71 | 3.38 | 2.81 | 2.50 | 78.85 | 77.20 | 58.10 | 5.84 | 6.10 |
| <i>SEm</i> ± | 0.04 | 0.004 | 0.005 | 0.01 | 0.005 | 0.05 | 0.07 | 0.21 | 0.17 | 0.005 | 0.003 |
| <i>LSD</i> ($P \leq 0.05$) | 0.14 | NS | 0.02 | 0.03 | NS | 0.16 | NS | 0.64 | 0.53 | NS | 0.011 |
| Interaction | NS | NS | NS | NS | NS | NS | NS | * | * | NS | NS |

CONCLUSIONS

The present study concluded that the application of gypsum at 52% GR gave maximum yield advantage, but it was at par with that of 23% GR. The study also revealed that the application of gypsum at 52% GR was highly ameliorative in terms of the physico-chemical properties of sodic soils. The grain quality of rice with the application of gypsum @ 52% GR value was superior over the treatments where no gypsum was applied, but it was at par with 23% GR Value.

ACKNOWLEDGEMENT

The author is thankful to Mr. Sanjeev Kumar Rai for the preparation of the manuscript and also thanks to Dr. Kedar Nath Rai, who has provided financial and valuable support during research works and publications.

REFERENCES

- Akhtar, J., Naseem, A., Mahmood, K., Nawaj, S., Quareshi, R.H. and Aslam, M.(2001) Response of selected wheat genotypes to salinity: growth and ionic relations. *Pakistan Journal of Soil Science* 19, 1-7.
- Al-Nabulsi, Y.A. (2001) Saline drainage water, irrigation frequency and crop species effects on some physical properties of soils. *Journal of Agronomy and Crop Science* 186, 15-20.
- Arora, Sanjay, Singh, Y.P., Mishra, V.K., Singh, A.K. and Sharma, D.K. (2015) Integrated effect of pressmud, chemical amendments and microbial bio-inoculant on sodic soil reclamation and productivity of rice-wheat. In *Advances in Soil and Water Resource Management for Food and Livelihood Security in Changing Climate* (S. Bhan and S. Arora, Eds.), SCSIR, New Delhi, pp. 519-528.
- Brugnoli, B. and Lauter, M. (1991) Effect of salinity on stomatal conductance, photosynthetic capacity and carbon isotope discrimination of salt-tolerant cotton and *Phaseolus vulgaris*. *Plant Physiology* 95, 628-635.
- Chhabra, R., Singh, N.T. and Hansen, A. (1989) Interaction between type of amendment and P availability. Annual Report. Central Soil Salinity Research Institute, Karnal, India, p.38.

Chhabra, R. and Abrol, I.P. (1977) Reclaiming effect of rice grown in sodic soils. *Soil Science* 124, 49-55.

Dutta, K.K., Mishra, B., Singh, R.K., Singh, K.N. and Singh, Y.P. (1996) Technological options for reclamation of salt affected soil under resource constraint environment in Uttar Pradesh. *Agricultural Situation in India*. April 1996, pp.17-21.

Ehert, D.L., Redman, R.E., Harvey, B.L. and Cipywnyk, A. (1990) Salinity induced calcium deficiency in wheat and barley. *Plant and Soil* 128, 143-151.

El-Hendawy, S.E., Hu, Y., Yakout, G.M., Awad, A.M., Hafiz, S.E. and Schmidhalter, U. (2005) Evaluating salt tolerance of wheat genotypes using multiple parameters. *European Journal of Agronomy* 22, 243- 253.

Flower, T.J., Garcia, A., Koyama, M. and Yeo, A.R. (1997) Breeding for salt tolerance in crop plants – the role of molecular biology. *Acta Physiologia Plantarum* 19, 427-433.

Flower, T.J., Hajbagheri, M.A. and Yeo, A.R. (1990) Ion accumulation in the cell wall of rice plants growing under saline conditions. *Plant Cell Environment* 14, 319-325.

Foss Analytical (2003) The determination of nitrogen according to Kjeldahl using block digestion and steam distillation. FOSS Analytical AB, Hoganas, Sweden. AN 300.

Houshmand, S., Arzani, A., Maibody, S.A.M. and Feizi, M. (2005) Evaluation of salt tolerance genotypes of durum wheat derived from in vitro and field experiment. *Field Crops Research* 91, 345-354.

Jackson, M.L. (1973) *Soil Chemical Analysis*. Prentice Hall of India Private Limited, New Delhi.

Jardat, A.A., Shahid, M. and Al-Maskri, A. (2004) Genetic diversity in the Batini barley landrace from Oman: II. Response to salinity stress. *Crop Science* 24, 310- 315.

K.C. (1994) IR 4630 derived lines are stable high yielder under saline conditions in Kerala, India, *IRRN* 19, 21.

Kadamhdad, G.R., Babhulkar, V.P. and Banhulkar, P.S. (1996) *Annals of Plant Physiology* 10, 153.

Kingsbury, R. and Epstein, E. (1986) Selection of salt-resistant spring wheat. *Crop Science* 24, 310-315.

Kumar, Ashok, Sharma, D.K. and Sharma, H.C. (1994) Growth, yield and water use efficiency of wheat as influenced by irrigation and nitrogen levels in sodic soils. *Indian Journal of Agronomy* 39, 220-224.

Marcel Dekker Inc., New York.

Meri, A. (1984) Plant response to salinity: Experimental methodology and application to the field soil salinity under irrigation process and management. Springer Verlag, Berlin.

Nitant, H.C. and Bhumbra, D.R. (1974) Transformation and movement of nitrogen fertilizers in sodic soils. Annual Report, Central Soil Salinity Research Institute, Karnal, India. p.24.

NRSA and Associates (1996) Mapping of salt-affected soils of India, 1: 250,000 map sheets, Legend. NRSA Hyderabad.

Obrejanu, G. and Sandhu, G.A. (1971) Amelioration of solonetz and solonetzized soils in the socialist Republic of Romania. In *European Solonetz Soils and their Reclamation* (Szabolcs, Ed.), Akademi Kiado, Budapest, pp. 99-130.

Pesarrakli, M. (1999) *Handbook of Plant and Crop Stress*.

Ponnamperuma, F.N. (1984) Role of cultivar tolerance in increasing rice production on salinized lands. In *Salinity Tolerance in Plants: Strategies for Crop Improvement* (R.C. Staples and G.H. Toenniessen, Eds.). Wiley, New York, pp. 255-271.

Rao, D.L.N. and Batra L. (1983) Ammonia volatilization from applied nitrogen in alkali soils. *Plant and Soil* 70, 219-228.

Richards, L.A. (1954) *Diagnosis and improvement of saline and alkali soils*. USDA (Washington) Handbook No. 60, pp. 112.

Saneoka, H., Shlota, K., Kurban, H., Chaudhary, M., Premchandra, G.S. and Fujita, K. (1999) Effect of salinity on growth and solute accumulation in two wheat lines differing in salt tolerance. *Soil Science Plant Nutrition* 45, 873-880.

Schoonover, W.R. (1952) Examination of soils for alkali. University of California, extension Service, Berkeley, California, USA.

Shylaraj, K.S., George, K.M., Shashidharan, N.K. and Nair,

Singh, G., Sharma, D.P. and Mehta, K.K. (1983) Effect of sources, levels and methods of nitrogen application on yield attributes, yield and N uptake of rice in a highly sodic soil. *46th Annual Convention of Indian Society of Soil Science*. Hisar, India.

Singh, Y.P., Singh, R. and Kumar, Neeraj (2008) Response of rice and wheat to Gypsum rates in sodic soils. *Indian Journal of Agricultural Sciences* 78, 362-365.

Singh, Y.P., Singh, R., Sharma, D.K., Nayak, A.K., Mishra, V.K. and Singh, G. (2009) Synergy of reduced gypsum and salt tolerant varieties: A low-cost sodic soil reclamation technology. *Technical Bulletin: CSSRI / Lucknow/2009/03*.

Sridhar, V., Soundarajan, M.S., Sudhakar Rao, R. and Stree Amula, C. (1985) Response of groundnut to rates, time and methods of gypsum application. *Madras Agricultural Journal* 72, 47-53.

Subbiah, B.V. and Asija, G.L. (1956) A rapid procedure for estimation of available nitrogen in soils. *Current Science* 25, 259-263.

Tanji, K.K. (1990) In *Agricultural Salinity Assessment and Management* (K.K. Tanji, Ed.), American Society of Civil Engineers, New York, pp. 619.

Yeo, A.R. and Flower, T.J. (1986) Salinity resistance in rice and a pyramiding approach to breeding varieties for saline soils. *Australian Journal of Plant Physiology* 13, 161-173.

Zapata, F.J., Alejar, M.S., Torrizo, L.B., Novero, A.U., Singh, V.P. and Senadhira, D. (1991) Field performance of another culture derived lines from F1 crosses of indica rice under saline and non-saline conditions. *Theoretical and Applied Genetics* 83, 6-11.

